

## Symptoms reported in SARS-CoV-2 PCR negatives

Elisabeth Dietz, Ann Sarah Walker on behalf of the COVID-19 Infection Survey


Nuffield Department of Medicine, University of Oxford

*Current Data Run: August 1<sup>st</sup> 2022 – November 28<sup>th</sup> 2022*


Generalized Additive Models (GAMs) were fitted on data from study assessments conducted in participants from England between August 1<sup>st</sup> 2022 and November 28<sup>th</sup> 2022. Only visits with a confirmed negative PCR test for SARS-CoV-2 were included. Models were estimated separately by age group and for the overall population. GAMs were modeled using a negative-binomial distribution with log link, modeled with thin plate splines (k=35), penalized based on the third derivative. No other explanatory variables than time (measured in number of days) were included in the models. Breakpoints are identified by the first day zero is no longer in the 99% credible interval for the second derivative (corresponding to a 99.5% probability of change in trend) (<https://www.medrxiv.org/content/10.1101/2022.09.14.22279931v1>). Symptoms reported refer to symptoms experienced in the last 7 days

*ILI-CDC: Fever and cough/sore throat*


*ILI-ECDC: At least one systemic symptom (fever, fatigue, headache and myalgia), and at least one respiratory symptom (cough, sore throat, and shortness of breath)*





Prevalence of ILI-ECDC amongst Covid-negatives, by age group and overall



Prevalence of Fever amongst Covid-negatives, by age group and overall





*This work contains statistical data from ONS which is Crown Copyright. The use of the ONS statistical data in this work does not imply the endorsement of the ONS in relation to the interpretation or analysis of the statistical data. This work uses research datasets which may not exactly reproduce National Statistics aggregates.*